



Carbonic Anhydrase Inhibitors for Glaucoma

Carbonic anhydrase inhibitors for glaucoma

Your ophthalmologist has recommended you use a type of medicine called a **carbonic anhydrase inhibitor**. This kind of medicine is helpful in treating glaucoma.

Glaucoma is a disease that affects your eye's optic nerve, possibly leading to blindness. The optic nerve connects your eye to your brain so you can see. Glaucoma usually happens when fluid builds up in the front portion of your eye. Pressure increases in your eye, damaging the optic nerve and eventually stealing your sight.

Carbonic anhydrase inhibitors work by reducing the amount of fluid your eye produces. That helps lower eye pressure, and, hopefully, saves your vision.

Your ophthalmologist may have you take this carbonic anhydrase inhibitor as an eye drop or by mouth as a pill.

Also, you may need to take this medication along with other glaucoma medicine.

Every dose, every day—it can save your sight!

It is extremely important to use your glaucoma medicine exactly as your ophthalmologist tells you to. That includes taking every dose, every day. If you do not do this, you could become blind.

As with any medication, glaucoma medicine can cause side effects. Possible side effects of carbonic anhydrase inhibitors include:

- stinging eyes
- red eyes
- a skin rash (especially in people who are allergic to sulfa drugs)
- changes in how things taste to you (especially with carbonated drinks)
- Upset stomach (nausea)
- feeling tired
- decreased energy
- increase in urination (with the pills)
- tingling around the mouth and fingertips (with the pills)

Side effects may go away after you take the medicine for a while. But if the side effects still bother you, call your ophthalmologist. He or she may be able to lower your dose or change your medicine. **Never suddenly quit taking your medicine unless your doctor tells you to.**

Also, remember to tell your other doctors that you take a carbonic anhydrase inhibitor medicine for glaucoma.

How to use eye drops

Putting drops in your eyes may seem hard at first. But with practice, it should get much easier. Here are steps to take to properly use eye drops:

- Wash your hands before you remove the cap of the eye drop bottle. Do not touch the tip of the dropper. This helps prevent the spread of germs.


- Tilt your head back slightly. Then pull your lower lid away from the eye to form a "pocket."



- Look up and let the eye drop fall into the pocket. Do not touch your eye, eyelid or lashes with the bottle tip. Close your eyes (but do not blink). Keep them closed for 1 minute.



- With your other hand, hold the dropper tip over your eyelid pocket. To keep your hand steady, you may want to rest it on your other hand or forehead.



- Before opening your eyes, wipe away any excess drops or tears with a tissue. Then open your eyes.



- If you take more than one type of eye drop medicine, wait 5 minutes between each medicine. This allows the eye to absorb each type of drop.
- Put the cap back on the bottle and store it in a safe place away from heat. Keep them out of reach of children and pets. Do not keep eye drop medicine near other drops bottles (ear drops, toe fungus drops, or Superglue, for example). People often mistakenly grab the wrong bottle.