




What is a macular pucker?

Macular pucker happens when wrinkles, creases or bulges form on your macula. The macula must lie flat against the back of your eye to work properly. When the macula wrinkles or bulges, your central vision is affected.

With macular pucker, things can look wavy, or you may have trouble seeing details. You might notice a gray or cloudy area in your central vision. You may even have a blank spot in your central vision. Macular pucker will not affect your peripheral (side) vision.



👁️ Eye Words to Know

Retina: Layer of nerve cells lining the back wall inside the eye. This layer senses light and sends signals to the brain so you can see.

Macula: Small but important area in the center of the retina. You need the macula to clearly see details of objects in front of you.

Vitreous: Clear, gel-like substance that fills the inside of your eye. The vitreous helps the eye maintain its shape and also helps send light to the retina.

What causes macular pucker?

Age is the most common cause of macular pucker. As you get older, the vitreous begins to shrink and pull away from the retina. Usually the vitreous pulls away with no problems. But sometimes the vitreous can stick to the retina. Scar tissue forms, causing the retina and macula to wrinkle or bulge.

Who is at risk for macular pucker?

Aging is the most common risk factor for macular pucker. People who have other eye problems may also get a macular pucker. These problems include:

- Vitreous detachment, where the eye's vitreous pulls away from the retina
- Torn or detached retina
- Swelling inside the eye

- Serious damage to the eye (from surgery or injury)
- Problems with blood vessels in the retina

How is macular pucker diagnosed?

Your ophthalmologist will put drops in your eye to dilate (widen) your pupil. This allows him or her to look through a special lens at the inside of your eye.

Then he or she will take pictures of your eye using optical coherence tomography (OCT). With OCT, a machine scans the back of your eye. This provides very detailed pictures of the retina and macula. Your ophthalmologist studies these pictures to check for problems.



How is macular pucker treated?

How you are treated depends on your symptoms.

If your symptoms are mild, you might not need any treatment. Instead, your ophthalmologist may change your glasses or contact lens prescription to improve your vision. You might also choose to wear bifocals when you are looking at something close. Eye drops, medicine, and laser surgery do not help vision if you have macular pucker.

If your symptoms are more serious, your ophthalmologist may recommend a surgery called vitrectomy. Your ophthalmologist will remove some of the vitreous and scar tissue on your macula. This flattens the macula, returning it to its proper position. It is likely your vision will slowly improve. However, your sight may not be as good as it was before macular pucker.

Macular pucker



Your ophthalmologist uses tiny instruments to remove the wrinkled tissue on the macula.

What are vitrectomy surgery risks?

Like all surgery, vitrectomy has some risks. They include:

- Eye infection
- Bleeding in your eye
- A detached retina (where the retina lifts away from the back of the eye)
- Having the macular pucker happen again
- Cataract, when the lens in your eye becomes cloudy

Your ophthalmologist will talk about these risks and how vitrectomy surgery may help you.

Summary

Macular pucker happens when wrinkles or bulges form on your macula, affecting your central vision. In some cases, you won't need treatment. Instead, your ophthalmologist may change your glasses or contact lens prescription to improve your vision. If your symptoms are more serious, you may need vitrectomy surgery to flatten the macula and put it back in its proper position.

Get more information about macular puckers from EyeSmart—provided by the American Academy of Ophthalmology—at aao.org/macular-pucker-link.

COMPLIMENTS OF:

